

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
1 February 2001 (01.02.2001)

PCT

(10) International Publication Number  
**WO 01/07456 A1**

(51) International Patent Classification<sup>7</sup>: C07H 21/04, C12N 15/00, C12Q 1/68

(74) Agents: LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 08053 (US).

(21) International Application Number: PCT/US99/29589

(22) International Filing Date:  
14 December 1999 (14.12.1999)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/358,382 21 July 1999 (21.07.1999) US

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): ISIS PHARMACEUTICALS, INC. [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).

**Published:**

— With international search report.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): WARD, Donna, T. [US/US]; 11830 Caminito Ronaldo, Apartment 125, San Diego, CA 92128 (US). COWSERT, Lex, M. [US/US]; 3008 Newshire Street, Carlsbad, CA 92008 (US).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: ANTISENSE MODULATION OF JUN N-TERMINAL KINASE KINASE-1 EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of Jun N-terminal Kinase Kinase-1. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding Jun N-terminal Kinase Kinase-1. Methods of using these compounds for modulation of Jun N-terminal Kinase Kinase-1 expression and for treatment of diseases associated with expression of Jun N-terminal Kinase Kinase-1 are provided.

WO 01/07456 A1



-1-

**ANTISENSE MODULATION OF JUN N-TERMINAL KINASE KINASE-1  
EXPRESSION**

**FIELD OF THE INVENTION**

5       The present invention provides compositions and methods for modulating the expression of Jun N-terminal Kinase Kinase-1. In particular, this invention relates to antisense compounds, particularly oligonucleotides, specifically hybridizable with nucleic acids encoding human  
10 Jun N-terminal Kinase Kinase-1. Such oligonucleotides have been shown to modulate the expression of Jun N-terminal Kinase Kinase-1.

**BACKGROUND OF THE INVENTION**

15       One of the principal mechanisms by which cellular regulation is effected is through the transduction of extracellular signals across the membrane that in turn modulate biochemical pathways within the cell. Protein phosphorylation represents one course by which  
20 intracellular signals are propagated from molecule to molecule resulting finally in a cellular response. These signal transduction cascades are highly regulated and often overlapping as evidenced by the existence of many protein kinases as well as phosphatases. It is currently believed that a number of disease states and/or disorders are a  
25 result of either aberrant expression or functional mutations in the molecular components of kinase cascades. Consequently, considerable attention has been devoted to the characterization of these proteins.

30       Nearly all cell surface receptors use one or more of the mitogen-activated protein kinase (MAP kinase) cascades during signal transduction. Three distinct subgroups of the MAP kinases have been identified and each of these consists of a specific module of downstream kinases. One subgroup of the MAP kinases is the Jun N-terminal  
35 kinase/Stress activated protein kinase (JNK/SAPK) cascade.

-2-

This pathway was originally identified as an oncogene- and ultraviolet light-stimulated kinase pathway but is now known to be activated by growth factors, cytokines and T-cell costimulation (Moriguchi et al., *Adv. Pharmacol.*, 1996, 36, 121-137; Su and Karin, *Curr. Opin. Immunol.*, 1996, 8, 402-411).

Jun N-terminal Kinase Kinase-1 (also known as JNKK1, PRKMK4, MEK4, MKK4, SAPK/Erk kinase 1 and SEK1) is a dual-specificity MAP kinase kinase that acts to mediate signaling pathways initiated by cellular stressors (Derijard et al., *Science*, 1995, 267, 682-685). Jun N-terminal Kinase Kinase-1 has been shown to be regulated and activated to varying degrees by ultraviolet exposure (Butterfield et al., *Biochem. J.*, 1999, 338, 681-686), the  $\beta\gamma$  subunit of G-proteins (Yamauchi et al., *J. Biol. Chem.*, 1999, 274, 1957-1965) and mixed lineage kinase-2 (Cuenda and Dorow, *Biochem. J.*, 1998, 333, 11-15; Hirai et al., *J. Biol. Chem.*, 1998, 273, 7406-7412) suggesting that only certain stresses affect Jun N-terminal Kinase Kinase-1-mediated signaling pathways. Further investigations into the role of Jun N-terminal Kinase Kinase-1 in MAP kinase signaling cascades demonstrated that the interaction of Jun N-terminal Kinase Kinase-1 with both upstream and downstream components in the JNK signaling pathway is mediated through the amino terminus of the protein (Xia et al., *Genes Dev.*, 1998, 12, 3369-3381).

Three isoforms of Jun N-terminal Kinase Kinase-1 have been isolated and these are disclosed in US Patents numbers 5,736,381 and 5,804,427. Also disclosed are polynucleotides that encode three isoforms of Jun N-terminal Kinase Kinase-1, expression vectors comprising the polynucleotide sequence and host cells transformed with the recombinant expression vector as well as the Jun N-terminal Kinase



-3-

Kinase-1 polypeptide encoded by the polynucleotide (Davis et al., 1998; Davis et al., 1998).

Recently it was reported that Jun N-terminal Kinase Kinase-1 may act as a tumor suppressor gene, and sequence variants resulting from mutations and mismatches have been found in certain types of cancers including pancreatic, lung, breast and colon cancer cells (Su et al., *Cancer Res.*, 1998, 58, 2339-2342; Teng et al., *Cancer Res.*, 1997, 57, 4177-4182).

Disruption of the Jun N-terminal Kinase Kinase-1 gene in mice by homologous recombination resulted in embryos that displayed severe anemia and died by day 10.5. It was noted that, although hematopoiesis was normal, hepatogenesis and liver formation were severely impaired, with liver cells undergoing massive apoptosis and defects in AP-1 transcriptional activity (Ganiatsas et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1998, 95, 6881-6886; Nishina et al., *Development*, 1999, 126, 505-516; Yang et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1997, 94, 3004-3009).

Characterization of chimeric mice lacking Jun N-terminal Kinase Kinase-1 showed that mutation of Jun N-terminal Kinase Kinase-1 did not alter the induction of apoptosis in response to metabolic poisons but that T-cells failed to induce the expression of the death suppressor, Bcl-XL, suggesting that Jun N-terminal Kinase Kinase-1 is required to transduce cell-survival signals (Nishina et al., *J. Immunol.*, 1998, 161, 3416-3420).

To date, strategies aimed at inhibiting Jun N-terminal Kinase Kinase-1 function have involved the use of antibodies, dominant-negative forms of Jun N-terminal Kinase Kinase-1, gene knockouts in mice and SAPK pathway inhibitors. Disclosed in the PCT application WO 98/54203 are inhibitors of the SAPK pathway comprising antisense oligonucleotides to an upstream kinase, MEKK1, as well as

-4-

antisense oligonucleotides targeting the kinase isoforms that lie downstream of Jun N-terminal Kinase Kinase-1, SAPK1-3. Also disclosed are methods to inhibit the SAPK pathway using ribozymes, small molecule inhibitors and  
5 dominant-negative mutants (Mercola, 1998).

Currently, there are no known therapeutic agents which effectively inhibit the synthesis of Jun N-terminal Kinase Kinase-1 and consequently, there remains a long felt need for additional agents capable of specifically inhibiting  
10 Jun N-terminal Kinase Kinase-1 function.

Antisense technology is emerging as an effective means for reducing the expression of specific gene products and may therefore prove to be uniquely useful in a number of therapeutic, diagnostic, and research applications for the  
15 modulation of Jun N-terminal Kinase Kinase-1 expression.

#### SUMMARY OF THE INVENTION

The present invention is directed to antisense compounds, particularly oligonucleotides, which are targeted to a nucleic acid encoding Jun N-terminal Kinase  
20 Kinase-1, and which modulate the expression of Jun N-terminal Kinase Kinase-1. Pharmaceutical and other compositions comprising the antisense compounds of the invention are also provided. Further provided are methods of modulating the expression of Jun N-terminal Kinase  
25 Kinase-1 in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or  
30 condition associated with expression of Jun N-terminal Kinase Kinase-1 by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

#### DETAILED DESCRIPTION OF THE INVENTION

35 The present invention employs oligomeric antisense

-5-

compounds, particularly oligonucleotides, for use in modulating the function of nucleic acid molecules encoding Jun N-terminal Kinase Kinase-1, ultimately modulating the amount of Jun N-terminal Kinase Kinase-1 produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding Jun N-terminal Kinase Kinase-1. As used herein, the terms "target nucleic acid" and "nucleic acid encoding Jun N-terminal Kinase Kinase-1" encompass DNA encoding Jun N-terminal Kinase Kinase-1, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of Jun N-terminal Kinase Kinase-1. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with

-6-

the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding Jun N-terminal Kinase Kinase-1. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a

-7-

gene encoding Jun N-terminal Kinase Kinase-1, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself

-8-

as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as

5 "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target

10 regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It

15 has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently

20 complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick,

25 Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds.

"Complementary," as used herein, refers to the capacity for

30 precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to

35 each other at that position. The oligonucleotide and the

-9-

DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in

-10-

animals and man. Antisense oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 30 nucleobases (i.e. from about 8 to about 30 linked nucleosides). Particularly preferred antisense compounds are antisense oligonucleotides, even more preferably those comprising from about 12 to about 25 nucleobases. As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a



-11-

phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkyl-phosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and

-12-

those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

5        Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 10 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

15        Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short 20 chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; 25 methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts.

30        Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 35 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289;

-13-

5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

5 In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such  
10 oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular  
15 an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited  
20 to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

Most preferred embodiments of the invention are  
25 oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular  $-\text{CH}_2-\text{NH}-\text{O}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{O}-\text{CH}_2-$  [known as a methylene (methylimino) or MMI backbone],  $-\text{CH}_2-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-$ ,  $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{N}(\text{CH}_3)-\text{CH}_2-$  and  $-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-$  [wherein the  
30 native phosphodiester backbone is represented as  $-\text{O}-\text{P}-\text{O}-\text{CH}_2-$ ]  
] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S.  
35 patent 5,034,506.

-14-

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoethoxyethoxy, i.e., a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>3</sub>)<sub>2</sub>, also described in examples hereinbelow.

Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5'

-15-

terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar

5 structures include, but are not limited to, U.S.:

4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878;  
5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811;  
5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053;  
5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920,

10 certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or  
15 substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine  
20 (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil  
25 and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-  
30 methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of*  
35 *Polymer Science And Engineering*, pages 858-859, Kroschwitz,

-16-

J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-  
5 302, Crooke, S.T. and Lebleu, B. , ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines,  
10 including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds.,  
15 *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified  
20 nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121,  
25 5,596,091; 5,614,617; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and United States patent 5,750,692, which is commonly owned with the instant application and also herein incorporated by  
30 reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular

-17-

uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10, 1111-1118; Kabanov et al., *FEBS Lett.*, 1990, 259, 327-330; Svinarchuk et al., *Biochimie*, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654; Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277, 923-937).

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506;

-18-

5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241,  
5,391,723; 5,416,203, 5,451,463; 5,510,475; 5,512,667;  
5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481;  
5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and  
5 5,688,941, certain of which are commonly owned with the  
instant application, and each of which is herein  
incorporated by reference.

It is not necessary for all positions in a given  
compound to be uniformly modified, and in fact more than  
10 one of the aforementioned modifications may be incorporated  
in a single compound or even at a single nucleoside within  
an oligonucleotide. The present invention also includes  
antisense compounds which are chimeric compounds.  
"Chimeric" antisense compounds or "chimeras," in the  
15 context of this invention, are antisense compounds,  
particularly oligonucleotides, which contain two or more  
chemically distinct regions, each made up of at least one  
monomer unit, i.e., a nucleotide in the case of an  
oligonucleotide compound. These oligonucleotides typically  
20 contain at least one region wherein the oligonucleotide is  
modified so as to confer upon the oligonucleotide increased  
resistance to nuclease degradation, increased cellular  
uptake, and/or increased binding affinity for the target  
nucleic acid. An additional region of the oligonucleotide  
25 may serve as a substrate for enzymes capable of cleaving  
RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is  
a cellular endonuclease which cleaves the RNA strand of an  
RNA:DNA duplex. Activation of RNase H, therefore, results  
in cleavage of the RNA target, thereby greatly enhancing  
30 the efficiency of oligonucleotide inhibition of gene  
expression. Consequently, comparable results can often be  
obtained with shorter oligonucleotides when chimeric  
oligonucleotides are used, compared to phosphorothioate  
deoxyoligonucleotides hybridizing to the same target  
35 region. Cleavage of the RNA target can be routinely



-19-

detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be  
5 formed as composite structures of two or more  
oligonucleotides, modified oligonucleotides,  
oligonucleosides and/or oligonucleotide mimetics as  
described above. Such compounds have also been referred to  
in the art as hybrids or gapmers. Representative United  
10 States patents that teach the preparation of such hybrid  
structures include, but are not limited to, U.S.:  
5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878;  
5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355;  
5,652,356; and 5,700,922, certain of which are commonly  
15 owned with the instant application, and each of which is  
herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this  
invention may be conveniently and routinely made through  
the well-known technique of solid phase synthesis.  
20 Equipment for such synthesis is sold by several vendors  
including, for example, Applied Biosystems (Foster City,  
CA). Any other means for such synthesis known in the art  
may additionally or alternatively be employed. It is well  
known to use similar techniques to prepare oligonucleotides  
25 such as the phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are  
synthesized in vitro and do not include antisense  
compositions of biological origin, or genetic vector  
constructs designed to direct the in vivo synthesis of  
30 antisense molecules.

The compounds of the invention may also be admixed,  
encapsulated, conjugated or otherwise associated with other  
molecules, molecule structures or mixtures of compounds, as  
for example, liposomes, receptor targeted molecules, oral,  
35 rectal, topical or other formulations, for assisting in

-20-

uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S.:

5 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291;  
5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330;  
4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170;  
5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978;  
5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259;  
10 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of  
which is herein incorporated by reference.

The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to  
15 an animal including a human, is capable of providing  
(directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the  
disclosure is also drawn to prodrugs and pharmaceutically  
acceptable salts of the compounds of the invention,  
20 pharmaceutically acceptable salts of such prodrugs, and  
other bioequivalents.

The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof  
25 by the action of endogenous enzymes or other chemicals  
and/or conditions. In particular, prodrug versions of the  
oligonucleotides of the invention are prepared as SATE  
[(S-acetyl-2-thioethyl) phosphate] derivatives according to  
the methods disclosed in WO 93/24510 to Gosselin et al.,  
30 published December 9, 1993 or in WO 94/26764 to Imbach et  
al.

The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the  
35 desired biological activity of the parent compound and do

-21-

not impart undesired toxicological effects thereto.

Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid,

-22-

methylemaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 5 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, 10 methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfoic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, 15 N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable 20 cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of 25 pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, 30 hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, 35 tannic acid, palmitic acid, alginic acid, polyglutamic

-23-

acid, naphthalenesulfonic acid, methanesulfonic acid,  
p-toluenesulfonic acid, naphthalenedisulfonic acid,  
polygalacturonic acid, and the like; and (d) salts formed  
from elemental anions such as chlorine, bromine, and  
5 iodine.

The antisense compounds of the present invention can  
be utilized for diagnostics, therapeutics, prophylaxis and  
as research reagents and kits. For therapeutics, an  
animal, preferably a human, suspected of having a disease  
10 or disorder which can be treated by modulating the  
expression of Jun N-terminal Kinase Kinase-1 is treated by  
administering antisense compounds in accordance with this  
invention. The compounds of the invention can be utilized  
in pharmaceutical compositions by adding an effective  
15 amount of an antisense compound to a suitable  
pharmaceutically acceptable diluent or carrier. Use of the  
antisense compounds and methods of the invention may also  
be useful prophylactically, e.g., to prevent or delay  
infection, inflammation or tumor formation, for example.

20 The antisense compounds of the invention are useful  
for research and diagnostics, because these compounds  
hybridize to nucleic acids encoding Jun N-terminal Kinase  
Kinase-1, enabling sandwich and other assays to easily be  
constructed to exploit this fact. Hybridization of the  
25 antisense oligonucleotides of the invention with a nucleic  
acid encoding Jun N-terminal Kinase Kinase-1 can be  
detected by means known in the art. Such means may include  
conjugation of an enzyme to the oligonucleotide,  
radiolabelling of the oligonucleotide or any other suitable  
30 detection means. Kits using such detection means for  
detecting the level of Jun N-terminal Kinase Kinase-1 in a  
sample may also be prepared.

The present invention also includes pharmaceutical  
compositions and formulations which include the antisense  
35 compounds of the invention. The pharmaceutical

-24-

compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including  
5 ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral  
10 administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be  
15 particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical  
20 carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

Compositions and formulations for oral administration include powders or granules, suspensions or solutions in  
25 water or non-aqueous media, capsules, sachets or tablets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable.

Compositions and formulations for parenteral, intrathecal or intraventricular administration may include  
30 sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention  
35 include, but are not limited to, solutions, emulsions, and

-25-

liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

5       The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association  
10 the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the  
15 product.

      The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the  
20 present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The  
25 suspension may also contain stabilizers.

      In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams,  
30 jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may

-26-

be applied to the formulation of the compositions of the present invention.

#### Emulsions

5       The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter. (Idson, in *Pharmaceutical Dosage Forms*,  
10   Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman,  
15   Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases  
20   intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a  
25   water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to  
30   the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed.



-27-

Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex  
5 formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water  
10 stabilized in an oily continuous provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into  
15 the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing  
20 emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in  
25 *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of  
30 emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc.,  
35 New York, N.Y., 1988, volume 1, p. 199). Surfactants are

-28-

typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

-29-

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth),  
5 cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming  
10 strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides  
15 that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic  
20 acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing  
25 agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for  
30 their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of  
35 ease of formulation, efficacy from an absorption and

-30-

bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

10 In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar

-31-

heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

The phenomenological approach utilizing phase diagrams  
5 has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245;  
10 Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of  
15 thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers,  
20 polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol  
25 decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered  
30 film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to,  
35 water, an aqueous solution of the drug, glycerol, PEG300,

-32-

PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385; Ho et al., *J. Pharm. Sci.*, 1996, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the

-33-

gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

5       Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and  
10       nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et  
15       al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

#### Liposomes

20       There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and  
25       the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

      Liposomes are unilamellar or multilamellar vesicles  
30       which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as

-34-

efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each  
5 with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

Further advantages of liposomes include; liposomes  
10 obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical*  
15 *Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

20 Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As  
25 the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many  
30 drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the  
35 administered drug at the desired target, and the ability to



-35-

administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 1987, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 1992, 19, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine

-36-

(DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

5        Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, 1992, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

30        Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized

-37-

liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside  $G_{M1}$ , or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 1987, 223, 42; Wu et al., *Cancer Research*, 1993, 53, 3765). Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, 1987, 507, 64) reported the ability of monosialoganglioside  $G_{M1}$ , galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside  $G_{M1}$  or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C<sub>12</sub>15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with

-38-

polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klivanov et al. (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Patent Nos. 5,540,935 (Miyazaki et al.) and 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and

-39-

asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love et al. discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants

-40-

find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic  
5 esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block  
10 polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant  
15 is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and  
20 sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant  
25 is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

If the surfactant molecule has the ability to carry  
30 either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations  
35 and in emulsions has been reviewed (Rieger, in

-41-

*Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

#### Penetration Enhancers

5           In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually  
10   only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs  
15   across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

          Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-  
20   surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

25           Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and  
30   another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl  
35   ether) (Lee et al., *Critical Reviews in Therapeutic Drug*

-42-

Carrier Systems, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, 1988, 40, 252).

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C<sub>1-10</sub> alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

20

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium

30



-43-

glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate),  
5 ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydrofusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: *Remington's Pharmaceutical*  
10 *Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

15

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that  
20 absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal  
25 ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-  
30 methoxysalicylate and homovanilate), N-acyl derivatives of collagen, laureth-9 and N-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990,

-44-

7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

### Carriers

Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a

-45-

nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity *per se*) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., *Antisense Res. Dev.*, 1995, 5, 115-121; Takakura et al., *Antisense & Nucl. Acid Drug Dev.*, 1996, 6, 177-183).

20

#### Excipients

In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars,

30

-46-

microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

-47-

### Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at  
5 their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional  
10 materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere  
15 with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic  
20 pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for  
25 example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other  
30 chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include, but are not limited to, anticancer drugs such as daunorubicin, dactinomycin, doxorubicin, bleomycin, mitomycin, nitrogen mustard, chlorambucil, melphalan,  
35 cyclophosphamide, 6-mercaptopurine, 6-thioguanine,

-48-

cytarabine (CA), 5-fluorouracil (5-FU), floxuridine (5-FUdR), methotrexate (MTX), colchicine, vincristine, vinblastine, etoposide, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 1206-1228). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and

-49-

can generally be estimated based on  $EC_{50}$ s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

## EXAMPLES

### Example 1

#### Nucleoside Phosphoramidites for Oligonucleotide Synthesis Deoxy and 2'-alkoxy amidites

2'-Deoxy and 2'-methoxy beta-cyanoethyldiisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to

-50-

published methods [Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21, 3197-3203] using commercially available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

5 **2'-Fluoro amidites**

**2'-Fluorodeoxyadenosine amidites**

2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841] and United States patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S<sub>N</sub>2-displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

**2'-Fluorodeoxyguanosine**

The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP to give diisobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups.



-51-

Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

#### 2'-Fluorouridine

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

#### 10 2'-Fluorodeoxycytidine

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

#### 2'-O-(2-Methoxyethyl) modified amidites

2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

#### 20 2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]

5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum

-52-

was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

#### 2'-O-Methoxyethyl-5-methyluridine

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH<sub>3</sub>CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH<sub>2</sub>Cl<sub>2</sub>/acetone/MeOH (20:5:3) containing 0.5% Et<sub>3</sub>NH. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

#### 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70%

-53-

product. The solvent was evaporated and triturated with  $\text{CH}_3\text{CN}$  (200 mL). The residue was dissolved in  $\text{CHCl}_3$  (1.5 L) and extracted with 2x500 mL of saturated  $\text{NaHCO}_3$  and 2x500 mL of saturated  $\text{NaCl}$ . The organic phase was dried over  $\text{Na}_2\text{SO}_4$ ,  
5 filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with  $\text{EtOAc}$ /hexane/acetone (5:5:1) containing 0.5%  $\text{Et}_3\text{NH}$ . The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional  
10 was obtained from the impure fractions to give a total yield of 183 g (57%).

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine  
15 (106 g, 0.167 M),  $\text{DMF}$ /pyridine (750 mL of a 3:1 mixture prepared from 562 mL of  $\text{DMF}$  and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the  
20 addition of  $\text{MeOH}$ . Upon completion of the reaction, as judged by TLC,  $\text{MeOH}$  (50 mL) was added and the mixture evaporated at  $35^\circ\text{C}$ . The residue was dissolved in  $\text{CHCl}_3$  (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated  $\text{NaCl}$ . The water  
25 layers were back extracted with 200 mL of  $\text{CHCl}_3$ . The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product). The residue was purified on a 3.5 kg silica gel column and eluted using  $\text{EtOAc}$ /hexane(4:1). Pure product fractions were  
30 evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine**

A first solution was prepared by dissolving 3'-O-  
35 acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-

-54-

methylluridine (96 g, 0.144 M) in CH<sub>3</sub>CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH<sub>3</sub>CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POCl<sub>3</sub>,  
5 was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10°C, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored  
10 overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO<sub>3</sub> and 2x300 mL of saturated NaCl,  
15 dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M)  
20 in dioxane (500 mL) and NH<sub>4</sub>OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred  
25 to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH<sub>3</sub> gas was added and the vessel heated to 100°C for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with  
30 saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

35 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyl-

-55-

cytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl<sub>3</sub> (700 mL) and extracted with saturated NaHCO<sub>3</sub> (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO<sub>4</sub> and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et<sub>3</sub>NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite**

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra-(isopropyl)phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO<sub>3</sub> (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH<sub>2</sub>Cl<sub>2</sub> (300 mL), and the extracts were combined, dried over MgSO<sub>4</sub> and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

**2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminooxyethyl) nucleoside amidites**

**2'-(Dimethylaminooxyethoxy) nucleoside amidites**

2'-(Dimethylaminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(dimethylaminooxyethyl) nucleoside

-56-

amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected  
5 with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

**5'-O-tert-Butyldiphenylsilyl-0<sup>2</sup>-2'-anhydro-5-methyluridine**

0<sup>2</sup>-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese,  
10 Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one  
15 portion. The reaction was stirred for 16 h at ambient temperature. TLC (R<sub>f</sub> 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium  
20 bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to  
25 -10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

**5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine**  
30

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until

-57-

the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

**2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine**

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over P<sub>2</sub>O<sub>5</sub> under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL,

-58-

44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred  
5 for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-t-  
10 butyldiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine

2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-  
15 5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h the mixture was filtered, the filtrate was washed with ice cold CH<sub>2</sub>Cl<sub>2</sub> and the combined organic phase was washed with water, brine and  
20 dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under  
25 vacuum; residue chromatographed to get 5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine as white foam (1.95 g, 78%).

5'-O-tert-Butyldiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine

30 5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this



-59-

solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Aqueous NaHCO<sub>3</sub> solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO<sub>3</sub> (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

2'-O-(dimethylaminoxyethyl)-5-methyluridine

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%).

-60-

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine

2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%).

5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. Then the reaction mixture was dissolved in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-N,N,N<sup>1</sup>,N<sup>1</sup>-tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added. The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate 1:1). The solvent was evaporated, then the residue was dissolved in ethyl acetate (70mL) and washed with 5% aqueous NaHCO<sub>3</sub> (40mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

-61-

**2'-(Aminooxyethoxy) nucleoside amidites**

2'-(Aminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(aminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs.

- 5 Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

**N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

- 10 The 2'-O-aminooxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor  
15 amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinosso, C. J., WO 94/02501 A1 940203.) Standard protection procedures  
20 should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-  
25 5'-O-(4,4'-dimethoxytrityl)guanosine. As before the hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-  
30 dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

**2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites**

2'-dimethylaminoethoxyethoxy nucleoside amidites (also

-62-

known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, or 2'-DMAEOE nucleoside amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

5    2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine

2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. O<sup>2</sup>-, 2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

25        5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethyl  
aminoethoxy)ethyl]-5-methyl uridine

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x200 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers are washed with saturated NaHCO<sub>3</sub> solution, followed by saturated NaCl solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel

-63-

chromatography using MeOH:CH<sub>2</sub>Cl<sub>2</sub>:Et<sub>3</sub>N (20:1, v/v, with 1% triethylamine) gives the title compound.

5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite

Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the solvent evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

## Example 2

### Oligonucleotide synthesis

Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphinate oligonucleotides are prepared as described in U.S. Patent 5,508,270, herein incorporated by reference.

-64-

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

### Example 3

#### Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are

-65-

herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

5 Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

#### Example 4

##### PNA Synthesis

10 Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance  
15 with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

#### Example 5

##### Synthesis of Chimeric Oligonucleotides

Chimeric oligonucleotides, oligonucleosides or mixed  
20 oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of  
25 the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

30 [2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric

##### Phosphorothioate Oligonucleotides

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligo

-66-

nucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphor-  
 5 amidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl. The  
 10 fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect  
 15 all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25  
 20 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)] Chimeric Phosphorothioate

25 Oligonucleotides

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.  
 30

[2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy Phosphorothioate]--[2'-O-(2-Methoxyethyl) Phosphodiester] Chimeric Oligonucleotides

[2'-O-(2-methoxyethyl phosphodiester)]--[2'-deoxy phos-



-67-

phorothioate]--[2'-O-(methoxyethyl) phosphodiester]  
chimeric oligonucleotides are prepared as per the above  
procedure for the 2'-O-methyl chimeric oligonucleotide with  
the substitution of 2'-O-(methoxyethyl) amidites for the  
5 2'-O-methyl amidites, oxidization with iodine to generate  
the phosphodiester internucleotide linkages within the  
wing portions of the chimeric structures and sulfurization  
utilizing 3,4-dihydro-2H-benzodithiole-3-one 1,1 dioxide (Beaucage  
Reagent) to generate the phosphorothioate internucleotide  
10 linkages for the center gap.

Other chimeric oligonucleotides, chimeric oligonucleo-  
sides and mixed chimeric oligonucleotides/oligonucleosides  
are synthesized according to United States patent  
5,623,065, herein incorporated by reference.

#### 15 Example 6

##### Oligonucleotide Isolation

After cleavage from the controlled pore glass column  
(Applied Biosystems) and deblocking in concentrated  
ammonium hydroxide at 55°C for 18 hours, the  
20 oligonucleotides or oligonucleosides are purified by  
precipitation twice out of 0.5 M NaCl with 2.5 volumes  
ethanol. Synthesized oligonucleotides were analyzed by  
polyacrylamide gel electrophoresis on denaturing gels and  
judged to be at least 85% full length material. The  
25 relative amounts of phosphorothioate and phosphodiester  
linkages obtained in synthesis were periodically checked by  
<sup>31</sup>P nuclear magnetic resonance spectroscopy, and for some  
studies oligonucleotides were purified by HPLC, as  
described by Chiang et al., *J. Biol. Chem.* 1991, 266,  
30 18162-18171. Results obtained with HPLC-purified material  
were similar to those obtained with non-HPLC purified  
material.

-68-

**Example 7****Oligonucleotide Synthesis - 96 Well Plate Format**

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1 dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per known literature or patented methods. They are utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and deprotected with concentrated  $\text{NH}_4\text{OH}$  at elevated temperature (55-60°C) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

**Example 8****Oligonucleotide Analysis - 96 Well Plate Format**

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test

-69-

plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

5 **Example 9**

**Cell culture and oligonucleotide treatment**

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at  
10 measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following four cell types are provided for illustrative purposes, but other cell types can be routinely used.

15 **T-24 cells:**

The transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life  
20 Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and  
25 dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates  
30 and treated similarly, using appropriate volumes of medium and oligonucleotide.

**A549 cells:**

The human lung carcinoma cell line A549 was obtained  
35 from the American Type Culture Collection (ATCC) (Manassas,

-70-

VA). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

10 NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

HEK cells:

Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

Treatment with antisense compounds:

When cells reached 80% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200  $\mu$ L OPTI-MEM™-1 reduced-serum medium (Gibco BRL) and then treated with 130  $\mu$ L of OPTI-MEM™-1 containing 3.75  $\mu$ g/mL LIPOFECTIN™ (Gibco BRL) and the desired oligonucleotide at a final concentration of 150 nM. After 4 hours of treatment, the medium was

-71-

replaced with fresh medium. Cells were harvested 16 hours after oligonucleotide treatment.

#### Example 10

#### Analysis of oligonucleotide inhibition of Jun N-terminal

#### 5 Kinase Kinase-1 expression

Antisense modulation of Jun N-terminal Kinase Kinase-1 expression can be assayed in a variety of ways known in the art. For example, Jun N-terminal Kinase Kinase-1 mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions. Other methods of PCR are also known in the art.

Jun N-terminal Kinase Kinase-1 protein levels can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to Jun N-terminal Kinase Kinase-1 can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for

-72-

preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997. Preparation of monoclonal

5 antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F.M. et al.,  
10 *Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-  
15 10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

## 20 Example 11

### Poly(A)+ mRNA isolation

Poly(A)+ mRNA was isolated according to Miura et al., *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example,  
25 Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 60  $\mu$ L lysis buffer (10 mM  
30 Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55  $\mu$ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc.,

-73-

Irvine CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200  $\mu$ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate was then transferred to a fresh 96-well plate.

10 Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

#### Example 12

##### Total RNA Isolation

15 Total mRNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold  
20 PBS. 100  $\mu$ L Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100  $\mu$ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a  
25 QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15  
seconds. 1 mL of Buffer RPE was then added to each well of  
30 the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the  
35 QIAVAC™ manifold fitted with a collection tube rack

-74-

containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60  $\mu$ L water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60  $\mu$ L water.

5       The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are  
10       carried out.

#### Example 13

#### Real-time Quantitative PCR Analysis of Jun N-terminal Kinase Kinase-1 mRNA Levels

Quantitation of Jun N-terminal Kinase Kinase-1 mRNA  
15       levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput  
20       quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by  
25       including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE or FAM, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City,  
30       CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by  
35       the proximity of the 3' quencher dye. During



-75-

amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25  $\mu$ L PCR cocktail (1x TAQMAN™ buffer A, 5.5 mM MgCl<sub>2</sub>, 300  $\mu$ M each of dATP, dCTP and dGTP, 600  $\mu$ M of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLD™, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25  $\mu$ L poly(A) mRNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD™, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension). Jun N-terminal Kinase Kinase-1 probes and primers were designed to hybridize to the human Jun N-terminal Kinase Kinase-1 sequence, using published sequence information (GenBank accession number L36870, incorporated herein as SEQ ID NO:1).

-76-

For Jun N-terminal Kinase Kinase-1 the PCR primers were:

forward primer: CCACACATAGAGAGACTGAGAACACA (SEQ ID NO: 2)

reverse primer: TGCAGTGAAATCCCAGTGTTG (SEQ ID NO: 3) and

5 the PCR probe was: FAM-CATTGAGTCATCAGGAAACTGAAGATCTCCC-TAMRA

(SEQ ID NO: 4) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

10 For GAPDH the PCR primers were:

forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO: 5)

reverse primer: GAAGATGGTGATGGGATTTC (SEQ ID NO: 6) and the

PCR probe was: 5' JOE-CAAGCTTCCC GTTCTCAGCC- TAMRA 3' (SEQ ID NO: 7) where JOE (PE-Applied Biosystems, Foster City,

15 CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

#### Example 14

#### Northern blot analysis of Jun N-terminal Kinase Kinase-1 mRNA levels

20 Eighteen hours after antisense treatment, cell monolayers were washed twice with cold PBS and lysed in 1 mL RNazol™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA was prepared following manufacturer's recommended protocols. Twenty micrograms of total RNA was fractionated

25 by electrophoresis through 1.2% agarose gels containing 1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon, OH). RNA was transferred from the gel to HYBOND™-N+ nylon membranes (Amersham Pharmacia Biotech, Piscataway, NJ) by overnight capillary transfer using a

30 Northern/Southern Transfer buffer system (TEL-TEST "B" Inc., Friendswood, TX). RNA transfer was confirmed by UV visualization. Membranes were fixed by UV cross-linking using a STRATALINKER™ UV Crosslinker 2400 (Stratagene, Inc, La Jolla, CA).

-77-

Membranes were probed using QUICKHYB™ hybridization solution (Stratagene, La Jolla, CA) using manufacturer's recommendations for stringent conditions with a Jun N-terminal Kinase Kinase-1 specific probe prepared by PCR using the forward primer CCACACATAGAGAGACTGAGAACACA (SEQ ID NO: 2) and the reverse primer TGCAGTGAAATCCCAGTGTTG (SEQ ID NO: 3). To normalize for variations in loading and transfer efficiency membranes were stripped and probed for glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA). Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

#### Example 15

##### **Antisense inhibition of Jun N-terminal Kinase Kinase-1 expression- phosphorothioate oligodeoxynucleotides**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human Jun N-terminal Kinase Kinase-1 RNA, using published sequences (GenBank accession number L36870, incorporated herein as SEQ ID NO: 1). The oligonucleotides are shown in Table 1. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. L36870), to which the oligonucleotide binds. All compounds in Table 1 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. The compounds were analyzed for effect on Jun N-terminal Kinase Kinase-1 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

-78-

Table 1

Inhibition of Jun N-terminal Kinase Kinase-1 mRNA levels by  
phosphorothioate oligodeoxynucleotides

5	ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
	101249	Start Codon	1	ggagccgccattgttgggag	9	8
	101250	Start Codon	6	ggctcggagccgccattgtt	44	9
	101251	Start Codon	11	gctcgggctcggagccgcca	53	10
10	101252	Coding	25	gagccgccgccgctcgg	0	11
	101253	Coding	100	atgctgctgacggccgggtg	24	12
	101254	Coding	128	caacttcagtgtttgcgtt	58	13
	101255	Coding	171	gagtaaaccttgctgtagat	27	14
	101256	Coding	218	cagtctctctatgtgtgggt	31	15
15	101257	Coding	283	gtgaaatcccagtggtgttc	9	16
	101258	Coding	459	cactactccgcattactaca	44	17
	101259	Coding	614	gattttgcctaaaatttctt	0	18
	101260	Coding	718	atatttccacttctgtccag	33	19
	101261	Coding	1177	atgggagagctgggagtagc	0	20
20	101262	Coding	1188	aatcgacatacatgggagag	45	21
	101263	Stop Codon	1194	gatatcaatcgacatacatg	17	22
	101264	Stop Codon	1201	tagcagcgatatcaatcgac	45	23
	101265	Stop Codon	1206	tgatgtagcagcgatatcaa	28	24
	101266	Stop Codon	1209	gtctgatgtagcagcgatat	47	25
25	101267	3' UTR	1213	tagagtctgatgtagcagcg	21	26
	101268	3' UTR	1313	aacaccaatcttattgcaca	2	27
	101269	3' UTR	1361	aaggatgcacgttctaggtg	45	28
	101271	3' UTR	1431	gttcatcacaaaagagcagg	11	29
	101273	3' UTR	1505	tagaaatgaatttaagatgt	0	30
30	101275	3' UTR	1574	tgtgtccttccaggatttgg	30	31
	101277	3' UTR	1636	gctgtgcaagctcttctcac	37	32
	101279	3' UTR	1721	tatcacatcagtaattctatt	0	33
	101281	3' UTR	1778	aaatttggctgagcacatcg	47	34
	101282	3' UTR	1862	tattttaatgttcacctctt	0	35
35	101284	3' UTR	1953	caccttctttgacaagtggc	58	36
	101286	3' UTR	2122	gaaatatatagctgctacaa	0	37
	101289	3' UTR	2279	agacctggaccagtgatggc	40	38
	101290	3' UTR	2448	tttaaactagcgattatg	0	39

-79-

	101292	3' UTR	2532	gatcctcatctggtccaatg	40	40
	101294	3' UTR	2662	gcctcttgtaacagtgtga	53	41
	101296	3' UTR	2783	accactcacagtgtgtcttc	32	42
	101298	3' UTR	2873	ggagacacagcaatacgtgc	39	43
5	101300	3' UTR	2974	gagataatcacttttgaatt	0	44
	101301	3' UTR	3101	atggaggatgatggcttgct	17	45
	101302	3' UTR	3243	cacacaatagaacggtcaga	10	46
	101303	3' UTR	3347	tttgttttaataaagataca	12	47

10 As shown in Table 1, SEQ ID NOs 9, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 25, 26, 28, 31, 32, 34, 36, 38, 40, 41, 42 and 43 demonstrated at least 20% inhibition of Jun N-terminal Kinase Kinase-1 expression in this assay and are therefore preferred.

15 **Example 16:**

**Antisense inhibition of Jun N-terminal Kinase Kinase-1 expression- phosphorothioate 2'-MOE gapmer oligonucleotides**

In accordance with the present invention, a second series of oligonucleotides targeted to human Jun N-terminal Kinase Kinase-1 were synthesized. The oligonucleotide sequences are shown in Table 2. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. L36870), to which the oligonucleotide binds.

25 All compounds in Table 2 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. Cytidine residues in the 2'-MOE wings are 5-methylcytidines.

35 Data were obtained by real-time quantitative PCR as described in other examples herein and are averaged from two experiments. If present, "N.D." indicates "no data".

-80-

Table 2

Inhibition of Jun N-terminal Kinase Kinase-1 mRNA levels by  
chimeric phosphorothioate oligonucleotides having 2'-MOE  
wings and a deoxy gap

5

	ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
	101304	Start Codon	1	ggagccgccattggtgggag	66	8
	101305	Start Codon	6	ggctcggagccgccattggt	70	9
10	101306	Start Codon	11	gctcgggctcggagccgcca	76	10
	101307	Coding	25	gagccgccgccgccgctcgg	29	11
	101308	Coding	100	atgctgctgacggccgggtg	71	12
	101309	Coding	128	caacttcagtgtcttgcgtt	82	13
	101310	Coding	171	gagtaaacccttgctgtagat	53	14
15	101311	Coding	218	cagtctctctatgtgtgggt	60	15
	101312	Coding	283	gtgaaatcccagtggtgttc	56	16
	101313	Coding	459	cactactccgcattactaca	19	17
	101314	Coding	614	gattttgcctaaaaatttctt	11	18
	101315	Coding	718	atatttccacttctgtccag	40	19
20	101316	Coding	1177	atgggagagctgggagtagc	0	20
	101317	Coding	1188	aatcgacatacatgggagag	62	21
	101318	Stop Codon	1194	gatatcaatcgacatacatg	65	22
	101319	Stop Codon	1201	tagcagcgatatcaatcgac	72	23
	101320	Stop Codon	1206	tgatgtagcagcgatatcaa	55	24
25	101321	Stop Codon	1209	gtctgatgtagcagcgatat	70	25
	101322	3' UTR	1213	tagagtctgatgtagcagcg	64	26
	101323	3' UTR	1313	aacaccaatcttattgcaca	45	27
	101324	3' UTR	1361	aaggatgcacgttctaggtg	76	28
	101325	3' UTR	1431	gttcatcacaaaagagcagg	54	29
30	101326	3' UTR	1505	tagaaatgaatttaagatgt	20	30
	101327	3' UTR	1574	tgtgtccttccaggatttgg	44	31
	101328	3' UTR	1636	gctgtgcaagctcttctcac	69	32
	101329	3' UTR	1721	tatcacatcagtaattctatt	50	33
	101330	3' UTR	1778	aaatttggtgagcacatcg	48	34
35	101331	3' UTR	1862	tattttaatgttcacctctt	47	35
	101332	3' UTR	1953	caccttctttgacaagtggc	66	36
	101333	3' UTR	2122	gaaatatatagctgctacaa	41	37

-81-

	101334	3' UTR	2279	agacctggaccagtgatggc	59	38
	101335	3' UTR	2448	tttaaactagcgattatg	22	39
	101336	3' UTR	2532	gacctcatctggccaatg	5	40
	101337	3' UTR	2662	gcctcttgtaacagtgctga	80	41
5	101338	3' UTR	2783	accactcacagtgctgcttc	69	42
	101339	3' UTR	2873	ggagacacagcaatacgtgc	58	43
	101340	3' UTR	2974	gagataatcacttttgaatt	12	44
	101341	3' UTR	3101	atggaggatgatggcttgct	56	45
	101342	3' UTR	3243	cacacaatagaacggtcaga	42	46
10	101343	3' UTR	3347	ttgttttaataaagataca	11	47

As shown in Table 2, SEQ ID NOs 8, 9, 10, 12, 13, 14, 15, 16, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 31, 32, 33, 34, 35, 36, 37, 38, 41, 42, 43, 45 and 46 demonstrated at least 30% inhibition of Jun N-terminal Kinase Kinase-1 expression in this experiment and are therefore preferred.

#### Example 17

#### Western blot analysis of Jun N-terminal Kinase Kinase-1 protein levels

Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 ul/well), boiled for 5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to Jun N-terminal Kinase Kinase-1 is used, with a radiolabelled or fluorescently labeled secondary antibody directed against the primary antibody species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

-82-

What is claimed is:

1. An antisense compound 8 to 30 nucleobases in length targeted to a nucleic acid molecule encoding human Jun N-terminal Kinase Kinase-1, wherein said antisense compound  
5 specifically hybridizes with and inhibits the expression of human Jun N-terminal Kinase Kinase-1.

2. The antisense compound of claim 1 which is an antisense oligonucleotide.

3. The antisense compound of claim 2 wherein the  
10 antisense oligonucleotide has a sequence comprising SEQ ID NO: 9, 10, 12, 13, 14, 15, 17, 19, 21, 23, 24, 25, 26, 28, 31, 32, 34, 36, 38, 40, 41, 42, 43, 8, 16, 22, 27, 29, 33, 35, 37, 45 or 46.

4. The antisense compound of claim 2 wherein the  
15 antisense oligonucleotide has a sequence comprising SEQ ID NO: 9, 10, 12, 13, 14, 15, 19, 21, 23, 24, 25, 26, 28, 31, 32, 34, 36, 38, 41, 42 or 43.

5. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified  
20 internucleoside linkage.

6. The antisense compound of claim 5 wherein the modified internucleoside linkage is a phosphorothioate linkage.

7. The antisense compound of claim 2 wherein the  
25 antisense oligonucleotide comprises at least one modified sugar moiety.

8. The antisense compound of claim 7 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

9. The antisense compound of claim 2 wherein the  
30 antisense oligonucleotide comprises at least one modified nucleobase.

10. The antisense compound of claim 9 wherein the modified nucleobase is a 5-methylcytosine.



-83-

11. The antisense compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.

12. A composition comprising the antisense compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

13. The composition of claim 12 further comprising a  
5 colloidal dispersion system.

14. The composition of claim 12 wherein the antisense compound is an antisense oligonucleotide.

15. A method of inhibiting the expression of Jun N-terminal Kinase Kinase-1 in human cells or tissues comprising  
10 contacting said cells or tissues with the antisense compound of claim 1 so that expression of Jun N-terminal Kinase Kinase-1 is inhibited.

16. A method of treating a human having a disease or condition associated with Jun N-terminal Kinase Kinase-1  
15 comprising administering to said animal a therapeutically or prophylactically effective amount of the antisense compound of claim 1 so that expression of Jun N-terminal Kinase Kinase-1 is inhibited.

**THIS PAGE BLANK (USPTO)**

## SEQUENCE LISTING

<110> Donna T. Ward  
Lex M. Cowser  
ISIS PHARMACEUTICALS, INC.

<120> ANTISENSE MODULATION OF JUN N-TERMINAL KINASE KINASE-1 EXPRESSION

<130> RTSP-0034

<150> US 09/358,382

<151> 1999-07-21

<160> 47

<210> 1

<211> 3576

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (10)..(1209)

<400> 1

ctcccaaca atg gcg gct ccg agc ccg agc ggc ggc ggc ggc tcc ggg 48  
Met Ala Ala Pro Ser Pro Ser Gly Gly Gly Ser Gly  
1 5 10

ggc ggc agc ggc agc ggc acc ccc ggc ccc gta ggg tcc ccg gcg cca 96  
Gly Gly Ser Gly Ser Gly Thr Pro Gly Pro Val Gly Ser Pro Ala Pro  
15 20 25

ggc cac ccg gcc gtc agc agc atg cag ggt aaa cgc aaa gca ctg aag 144  
Gly His Pro Ala Val Ser Ser Met Gln Gly Lys Arg Lys Ala Leu Lys  
30 35 40 45

ttg aat ttt gca aat cca cct ttc aaa tct aca gca agg ttt act ctg 192  
Leu Asn Phe Ala Asn Pro Pro Phe Lys Ser Thr Ala Arg Phe Thr Leu  
50 55 60

aat ccc aat cct aca gga gtt caa aac cca cac ata gag aga ctg aga 240  
Asn Pro Asn Pro Thr Gly Val Gln Asn Pro His Ile Glu Arg Leu Arg  
65 70 75

aca cac agc att gag tca tca gga aaa ctg aag atc tcc cct gaa caa 288  
Thr His Ser Ile Glu Ser Ser Gly Lys Leu Lys Ile Ser Pro Glu Gln  
80 85 90

cac tgg gat ttc act gca gag gac ttg aaa gac ctt gga gaa att gga 336  
His Trp Asp Phe Thr Ala Glu Asp Leu Lys Asp Leu Gly Glu Ile Gly  
95 100 105

cga gga gct tat ggt tct gtc aac aaa atg gtc cac aaa cca agt ggg 384  
Arg Gly Ala Tyr Gly Ser Val Asn Lys Met Val His Lys Pro Ser Gly  
110 115 120 125

caa ata atg gca gtt aaa aga att cgg tca aca gtg gat gaa aaa gaa 432  
Gln Ile Met Ala Val Lys Arg Ile Arg Ser Thr Val Asp Glu Lys Glu

**THIS PAGE BLANK (USPTO)**

caa aaa caa ctt ctt atg gat ttg gat gta gta atg cgg agt agt gat Gln Lys Gln Leu Leu Met Asp Leu Asp Val Val Met Arg Ser Ser Asp 145 150 155	480
tgc cca tac att gtt cag ttt tat ggt gca ctc ttc aga gag ggt gac Cys Pro Tyr Ile Val Gln Phe Tyr Gly Ala Leu Phe Arg Glu Gly Asp 160 165 170	528
tgt tgg atc tgt atg gaa ctc atg tct acc tcg ttt gat aag ttt tac Cys Trp Ile Cys Met Glu Leu Met Ser Thr Ser Phe Asp Lys Phe Tyr 175 180 185	576
aaa tat gta tat agt gta tta gat gat gtt att cca gaa gaa att tta Lys Tyr Val Tyr Ser Val Leu Asp Asp Val Ile Pro Glu Glu Ile Leu 190 195 200 205	624
ggc aaa atc act tta gca act gtg aaa gca cta aac cac tta aaa gaa Gly Lys Ile Thr Leu Ala Thr Val Lys Ala Leu Asn His Leu Lys Glu 210 215 220	672
aac ttg aaa att att cac aga gat atc aaa cct tcc aat att ctt ctg Asn Leu Lys Ile Ile His Arg Asp Ile Lys Pro Ser Asn Ile Leu Leu 225 230 235	720
gac aga agt gga aat att aag ctc tgt gac ttc ggc atc agt gga cag Asp Arg Ser Gly Asn Ile Lys Leu Cys Asp Phe Gly Ile Ser Gly Gln 240 245 250	768
ctt gtg gac tct att gcc aag aca aga gat gct ggc tgt agg cca tac Leu Val Asp Ser Ile Ala Lys Thr Arg Asp Ala Gly Cys Arg Pro Tyr 255 260 265	816
atg gca cct gaa aga ata gac cca agc gca tca cga caa gga tat gat Met Ala Pro Glu Arg Ile Asp Pro Ser Ala Ser Arg Gln Gly Tyr Asp 270 275 280 285	864
gtc cgc tct gat gtc tgg agt ttg ggg atc aca ttg tat gag ttg gcc Val Arg Ser Asp Val Trp Ser Leu Gly Ile Thr Leu Tyr Glu Leu Ala 290 295 300	912
aca ggc cga ttt cct tat cca aag tgg aat agt gta ttt gat caa cta Thr Gly Arg Phe Pro Tyr Pro Lys Trp Asn Ser Val Phe Asp Gln Leu 305 310 315	960
aca caa gtc gtg aaa gga gat cct ccg cag ctg agt aat tct gag gaa Thr Gln Val Val Lys Gly Asp Pro Pro Gln Leu Ser Asn Ser Glu Glu 320 325 330	1008
agg gaa ttc tcc ccg agt ttc atc aac ttt gtc aac ttg tgc ctt acg Arg Glu Phe Ser Pro Ser Phe Ile Asn Phe Val Asn Leu Cys Leu Thr 335 340 345	1056
aag gat gaa tcc aaa agg cca aag tat aaa gag ctt ctg aaa cat ccc Lys Asp Glu Ser Lys Arg Pro Lys Tyr Lys Glu Leu Leu Lys His Pro 350 355 360 365	1104
ttt att ttg atg tat gaa gaa cgt gcc gtt gag gtc gca tgc tat gtt Phe Ile Leu Met Tyr Glu Glu Arg Ala Val Glu Val Ala Cys Tyr Val 370 375 380	1152

**THIS PAGE BLANK (USPTO)**

tgt aaa atc ctg gat caa atg cca gct act ccc agc tct ccc atg tat 1200  
 Cys Lys Ile Leu Asp Gln Met Pro Ala Thr Pro Ser Ser Pro Met Tyr  
 385 390 395

gtc gat tga tatcgctgct acatcagact ctagaaaaaa gggctgagag gaagcaagac 1259  
 Val Asp

gtaaagaatt ttcattcccgt atcacagtgt ttttattgct cgcccagaca ccatgtgcaa 1319  
 taagattggt gttcgtttcc atcatgtctg tatactcctg tcacctagaa cgtgcatcct 1379  
 tgtaatacct gattgatcac acagtgttag tgctggtcag agagacctca tcctgctctt 1439  
 ttgtgatgaa catattcatg aaatgtggaa gtcagtacga tcaagttggt gactgtgatt 1499  
 agatcacatc ttaaattcat ttctagactc aaaacctgga gatgcagcta ctggaatggt 1559  
 gttttgtcag acttccaaat cctggaagga cacagtgatg aatgtactat atctgaacat 1619  
 agaaactcgg gcttgagtga gaagagcttg cacagccaac gagacacatt gccttctgga 1679  
 gctgggagac aaaggaggaa tttactttct tcaccaagtg caatagatta ctgatgtgat 1739  
 attctgttgc tttacagtta cagttgatgt ttggggatcg atgtgctcag ccaaatttcc 1799  
 tgtttgaaat atcatgttaa attagaatga atttatcttt accaaaaacc atgttgcggt 1859  
 caaagagggtg aacattaaaa tatagagaca ggacagaatg tgttcttttc tcctctacca 1919  
 gtccatatttt tcaatgggaa gactcaggag tctgccactt gtcaaagaag gtgctgatcc 1979  
 taagaatttt tcattctcag aattcgggtg gctgccact tgatgttcca cctgccacaa 2039  
 accaccagga ctgaaagaag aaaacagtac agaaggcaaa gtttacagat gtttttaatt 2099  
 ctagtatttt atctggaaca acttgtagca gctatatatt tccccctggg cccaagcctg 2159  
 atacttttagc catcataact cactaacagg gagaagtagc tagtagcaat gtgccttgat 2219  
 tgattagata aagattttcta gtaggcagca aaagaccaa tctcagttgt ttgcttcttg 2279  
 ccatcactgg tccaggtctt cagtttccga atctctttcc cttccccctgt ggtctattgt 2339  
 cgctatgtga cttgcgctta atccaatatt ttgccttttt tctatatcaa aaaaccttta 2399  
 cagttagcag ggatgttcct taccgaggat ttttaacccc caatctctca taatcgctag 2459  
 tgtttaaaag gctaagaata gtggggccca accgatgtgg taggtgataa agaggcatct 2519  
 tttctagaga cacattggac cagatgagga tccgaaacgg cagcctttac gttcatcacc 2579  
 tgctagaacc tctcgtagtc catcaccatt tcttggcatt ggaattctac tggaaaaaaa 2639  
 tacaaaaagc aaaacaaaac cctcagcact gttacaagag gccatttaag tatcttgtgc 2699  
 ttcttcactt acccattagc caggtttctca ttaggttttg cttgggcctc cctggcactg 2759  
 aaccttaggc tttgtatgac agtgaagcag cactgtgagt ggttcaagca cactggaata 2819

**THIS PAGE BLANK (USPTO)**



taaaacagtc atggcctgag atgcagggtga tgccattaca gaaccaaadc gtggcacgta 2879  
 ttgctgtgtc tcctctcaga gtgacagtca taaatactgt caaacaataa agggagaatg 2939  
 gtgctgttta aagtcacatc cctgtaaatt gcagaattca aaagtgatta tctctttgat 2999  
 ctacttgctt catttcccta tcttctcccc cacggatatcc taaacttttag acttcccact 3059  
 gttctgaaag gagacattgc tctatgtctg ccttcgacca cagcaagcca tcctcctcca 3119  
 ttgctccccg ggactcaaga ggaatctgtt tctctgctgt caacttccca tctggctcag 3179  
 cataggggtca ctttgccatt atgcaaatgg agataaaagc aattctggct gtccaggagc 3239  
 taatctgacc gttctattgt gtggatgacc acataagaag gcaatttttag tgtattaatc 3299  
 atagattatt ataaactata aacttaaggg caaggagttt attacaatgt atctttatta 3359  
 aaacaaaagg gtgtatagtg ttcacaaact gtgaaaatag tgtaagaact gtacattgtg 3419  
 agctctgggtt atttttctct tgtaccatag aaaaatgtat aaaaattatc aaaaagctaa 3479  
 tgtgcaggga tattgcctta tttgtctgta aaaaatggag ctgagtaaca taactgcttc 3539  
 ttggagcttt ggaatatttt atcctgtatt cttgtttt 3576

<210> 2  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<223> PCR Primer

<400> 2  
 ccacacatag agagactgag aacaca 26

<210> 3  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<223> PCR Primer

<400> 3  
 tgcagtgaag tcccagtggt g 21

<210> 4  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence

<223> PCR Probe

<400> 4  
 cattgagtca tcaggaaaac tgaagatctc cc 32

**THIS PAGE BLANK (USPTO)**

<210> 5  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<223> PCR Primer

<400> 5  
gaaggtgaag gtcggagtc 19

<210> 6  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> PCR Primer

<400> 6  
gaagatggtg atgggatttc 20

<210> 7  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> PCR Probe

<400> 7  
caagcttccc gttctcagcc 20

<210> 8  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 8  
ggagccgcca ttggtgggag 20

<210> 9  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 9  
ggctcggagc cgccattgtt 20

<210> 10  
<211> 20  
<212> DNA  
<213> Artificial Sequence

**THIS PAGE BLANK (USPTO)**

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 10

gctcgggctc ggagccgcca

20

&lt;210&gt; 11

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 11

gagccgcccgc cgccgctcgg

20

&lt;210&gt; 12

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 12

atgctgctga cgccggggtg

20

&lt;210&gt; 13

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 13

caacttcagt gctttgcgtt

20

&lt;210&gt; 14

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 14

gagtaaacct tgctgtagat

20

&lt;210&gt; 15

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 15

cagtctctct atgtgtgggt

20

**THIS PAGE BLANK (USPTO)**

<210> 16  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 16  
gtgaaatccc agtggtgttc 20

<210> 17  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 17  
cactactccg cattactaca 20

<210> 18  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 18  
gattttgcct aaaatttctt 20

<210> 19  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 19  
atatttccac ttctgtccag 20

<210> 20  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide

<400> 20  
atgggagagc tgggagtagc 20

<210> 21  
<211> 20  
<212> DNA

**THIS PAGE BLANK (USPTO)**



<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 21

aatcgacata catgggagag

20

<210> 22

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 22

gatatcaatc gacatacatg

20

<210> 23

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 23

tagcagcgat atcaatcgac

20

<210> 24

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 24

tgatgtagca gcgatatcaa

20

<210> 25

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 25

gtctgatgta gcagcgatat

20

<210> 26

<211> 20

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

**THIS PAGE BLANK (USPTO)**

<400> 26  
tagagtctga tgtagcagcg

20

<210> 27  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 27  
aacaccaatc ttattgcaca

20

<210> 28  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 28  
aaggatgcac gttctaggtg

20

<210> 29  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 29  
gttcatcaca aaagagcagg

20

<210> 30  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 30  
tagaaatgaa tttaagatgt

20

<210> 31  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 31  
tgtgtccttc caggatttgg

20

**THIS PAGE BLANK (USPTO)**

<210> 32  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 32  
gctgtgcaag ctcttctcac

20

<210> 33  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 33  
tatcacatca gtaatctatt

20

<210> 34  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 34  
aaatttggt gagcacatcg

20

<210> 35  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 35  
tattttaatg ttcacctctt

20

<210> 36  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 36  
caccttcttt gacaagtggc

20

<210> 37  
<211> 20  
<212> DNA  
<213> Artificial Sequence

**THIS PAGE BLANK (USPTO)**

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 37

gaaatatata gctgctacaa

20

&lt;210&gt; 38

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 38

agacctggac cagtgatggc

20

&lt;210&gt; 39

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 39

tttaaact agcgattatg

20

&lt;210&gt; 40

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 40

gacccatc tggccaatg

20

&lt;210&gt; 41

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 41

gcctcttga acagtgtga

20

&lt;210&gt; 42

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

## &lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 42

accactcaca gtgctgcttc

20

**THIS PAGE BLANK (USPTO)**



<210> 43  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 43  
ggagacacag caatacgtgc

20

<210> 44  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 44  
gagataatca cttttgaatt

20

<210> 45  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 45  
atggaggatg atggcttgct

20

<210> 46  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 46  
cacacaatag aacggtcaga

20

<210> 47  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 47  
tttgttttaa taaagataca

20

**THIS PAGE BLANK (USPTO)**

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 9589

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : C07H 21/04; C12N 15/00; C12Q 1/68

US CL : 435/6, 91.1, 91.31; 536/23.1, 24.1, 24.5

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 91.1, 91.31; 536/23.1, 24.1, 24.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
WEST, MEDLINE, BIOSIS, EMBASE, CAPLUS, SCISEARCH**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,736,381 A (DAVIS et al) 07 April 1998, see especially column 3.	1, 2, 5-15

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

•	Special categories of cited documents:	• T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
• A	document defining the general state of the art which is not considered to be of particular relevance		
• B	earlier document published on or after the international filing date	• X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an ... inventive step when the document is taken alone
• L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	• Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
• O	document referring to an oral disclosure, use, exhibition or other means		
• P	document published prior to the international filing date but later than the priority date claimed	• A	document member of the same patent family

Date of the actual completion of the international search

23 FEBRUARY 2000

Date of mailing of the international search report

14 MAR 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SEAN MCGARRY

Telephone No. (703) 308-0196

**THIS PAGE BLANK (USPTO)**

## INTERNATIONAL SEARCH REPORT

International application No.

US99/29589

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : C07H 21/04; C12N 15/00; C12Q 1/68

US CL : 435/6, 91.1, 91.31; 536/23.1, 24.1, 24.5

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 91.1, 91.31; 536/23.1, 24.1, 24.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST, MEDLINE, BIOSIS, EMBASE, CAPLUS, SCISEARCH

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,736,381 A (DAVIS et al) 07 April 1998, see especially column 3.	1, 2, 5-15

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*B\* earlier document published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*Z\* document member of the same patent family

Date of the actual completion of the international search

23 FEBRUARY 2000

Date of mailing of the international search report

14 MAR 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SEAN MCGARRY

Telephone No. (703) 308-0196

**THIS PAGE BLANK (USPTO)**